PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

	ssification 6:		(11) International Publication Number: WO 96/3720			
A61K 31/20 // (A61)	K 31/20, 31:20)	A1	(43) International Publication Date: 28 November 1996 (28.11.96			
21) International Application Number: PCT/GB96/01265 22) International Filing Date: 24 May 1996 (24.05.96) 30) Priority Data: 9510636.5 25 May 1995 (25.05.95) GB		CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG KP, KR, KZ, LK, LR, LT, LU, LV, MD, MG, MN, MW MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE LS, MW, SD, SZ, UG), European patent (AT, BE, CH, DE				
HOLDINGS PLC [G Meadows, Guildford, (72) Inventor; and (75) Inventor/Applicant (for Uline [GB/GB]; Weyve	Applicant (for all designated States except US): SCOTIA HOLDINGS PLC [GB/GB]; Efamol House, Woodbridge Meadows, Guildford, Surrey GU1 1BA (GB). Inventor; and Inventor/Applicant (for US only): STORDY, Barbara, Jacqueline [GB/GB]; Weyvern House, Weyvern Park, Portsmouth Road, Peasmarsh, Guildford, Surrey GU3 1NA (GB).		Published With international search report. Before the expiration of the time limit for amending claims and to be republished in the event of the receipt			
(74) Agents: STURT, Cliffor Bedford Row, Holbon	rd, Mark et al.; J. Miller & n, London WC1R 4JH (GB).		4			
	·					
	<u> </u>					
(54) Title: USE OF DHA AS	S A PHARMACEUTICAL C	OMPC	NOITIS			
(57) Abstract						
A method of combating manufacture of a medicament	t for such purposes, charact	erised	by administering/incorporating DHA as an active in effective amou			
A method of combating manufacture of a medicament	t for such purposes, charact	erised	by administering/incorporating DHA as an active in effective amount			
A method of combating manufacture of a medicament	t for such purposes, charact	erised	by administering/incorporating DHA as an active in effective amount			
A method of combating manufacture of a medicament	t for such purposes, charact	erised	by administering/incorporating DHA as an active in effective amount			
A method of combating manufacture of a medicament	t for such purposes, charact	erised	by administering/incorporating DHA as an active in effective amount			
A method of combating manufacture of a medicament	t for such purposes, charact	erised	by administering/incorporating DHA as an active in effective amou			
A method of combating manufacture of a medicament optionally in association with	t for such purposes, charact	erised	on or dark adaptation in dyslexics or normal persons, and a method by administering/incorporating DHA as an active in effective amou 6 essential fatty acids.			
A method of combating manufacture of a medicament	t for such purposes, charact	erised	by administering/incorporating DHA as an active in effective amou			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
ΑU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU .	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PŁ	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgystan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic	SD	Sudan
CF	Central African Republic		of Korea	SE	Sweden
CG	Congo	KR	Republic of Korea	SG	Singapore
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LR	Liberia	SZ	Swaziland
CS	Czechoslovakia	LT	Lithuania	TD	Chad
CZ	Czech Republic	LU	Luxembourg	TG	Togo
DE	Germany	LV	Latvia	TJ	Tajikistan
DK	Denmark	MC	Monaco	TT	Trinidad and Tobago
EE	Estonia	MD	Republic of Moldova	UA	Ukraine
KS	Spain	MG	Madagascar	UG	Uganda
FI	Finland	ML	Mali	US	United States of America
FR	France	MN	Mongolia	UZ.	Uzbekistan
GA	Gabon	MR	Mauritania	VN	Viet Nam

USE OF DHA AS A PHARMACEUTICAL COMPOSITION

Background

Dyslexia is a major problem of human development. It is a disorder manifest by difficulty in learning to read despite conventional instruction, adequate intelligence and socio-cultural opportunities, and arises from fundamental cognitive disabilities.

Dyslexia is four to five times commoner in boys than girls, commoner in children with atopic eczema and asthma than those without, and also commoner in offenders than the law abiding. It is associated with a loss of normal brain asymmetry which is demonstrable using modern brain scanning methods and also on functional tests. It is recognised to be a disorder with an organic basis.

Current Work

Docosahexaenoic (DHA) is a major constituent of the retina, of nerve tissue and of the brain, and studies have now been done to look for abnormalities in dyslexic individuals in the intake of nutrients known to be important in brain function, particularly DHA. The first study looked at 33 dyslexic children and 48 controls and was directed to maternal diet during pregnancy. Mothers were given a comprehensive questionnaire, directed particularly at the levels of alpha-linolenic acid (ALA) in the diet and the ratio of ALA to linoleic acid (LA). (ALA is converted to DHA within the body and that conversion can be inhibited by a high intake of LA). The pathways of metabolism of LA and ALA are shown in Figure 1. The study showed that the mothers of dyslexic children were significantly more likely to have consumed a diet with a low ALA:LA ratio during pregnancy.

A second study focused specifically on the intakes of foods, in particular fish and meats, which contain DHA itself. This study demonstrated that mothers of dyslexic children consumed diets lower in DHA.

SUBSTITUTE SHEET (RULE 26)

WO 96/37200 PCT/GB96/01265

It was therefore decided to investigate whether DHA supplementation might be beneficial in dyslexic individuals. A test capable of quickly demonstrating an effect was needed, and as DHA is known to be particularly important in the function of the retinal rods, required for vision in the dark, it was decided to test for reduced retinal DHA levels in dyslexia, as indicated by dark adaptation. Ten adults with dyslexia (4 females and 6 males) and ten control subjects (6 females and four males) were recruited. They were all young adults with age range of 18-26 years. Dark adaptation was tested using a standard instrument, the Friedman Visual Field Analyser, set for the dark adaptation function. One eye was occluded, bright light was shone in the other eye to bleach the retina and the room was darkened. Measurements of dark adaptation were made at one minute intervals by assessing the intensity of very brief flashes of light which could just be detected. Measurements were continued until no further adaptation was observed.

Because dark adaptation can be influenced by a number of known nutrients, including vitamins A and C, riboflavin, nicotinic acid, thiamin and zinc, all subjects were asked to keep a careful 7 day record of food eaten and the results analysed by the diet analysis program, COMPEAT 4. No differences between the control and dyslexia groups in any of these nutrients were detected.

The results for the two groups are presented in Figure 2 which shows the means and standard deviations. The dyslexics at every time point show poorer dark adaptation than the controls and the differences between the two groups are statistically significant (p < 0.05).

Tests for influence of DHA on dark adaptation was therefore conducted. For a period of one month 5 dyslexics and 5 controls were given 4 capsules per day of a fish oil which contained 120 mg of DHA per capsule, with no vitamin A or vitamin D. Dark adaptation was then retested.

In four of the controls, DHA had no effects on dark adaptation, although in one subject adaptation clearly improved. In contrast, in the dyslexic subjects as shown in Figure 3, DHA consistently and significantly improved dark adaptation (p < 0.041).

WO 96/37200 PCT/GB96/01265

Subsequently, DHA supplements given to dyslexic children have been found to be associated with apparent improvements in reading ability and behaviour. These reports are currently anecdotal and subjective but more formal controlled studies are in preparation.

The Invention

The invention lies in combating dyslexia or inadequate night vision or dark adaptation in dyslexics or normal individuals, by administering DHA or a precursor n-3 EFA, and particularly in:-

- 1. A method of treating the conditions by administering DHA or precursor to children and adults showing them.
- 2. A method of preventing the conditions by administering DHA or precursor to women during pregnancy and to infants in the year after birth.

The invention also lies in a method of preparation of a medicament, for use in combating dyslexia or inadequate night vision or dark adaptation as above, when DHA or precursor is used.

Although DHA is a key fatty acid in both the retina and the brain, the n-6 fatty acids derived from linoleic acid (Figure 1) are also important in these tissues. Because DHA and eicosapentaenoic acid (EPA) (which is usually associated with DHA in fish oils) can in some circumstances inhibit conversion of LA to gamma-linolenic acid (GLA), it may be appropriate in some situations to provide, with the DHA, supplements of LA or preferably GLA, dihomo-gamma-linolenic acid (DGLA) or arachidonic acid (AA) to prevent depletion of these important fatty acids. It may also be appropriate to provide other n-3 essential fatty acids such as alpha-linolenic acid, stearidonic acid (SA) or EPA for their specific properties rather than as DHA precursors.

The fatty acids may be delivered in any appropriate form which can raise the levels of DHA and/or the other fatty acids in the blood. Appropriate forms are the free fatty acids; their salts, including lithium salts; esters; diesters of 1,3-propane diol and other

SUBSTITUTE SHEET (RULE 26)

diols; amides; alcohols; tri-, di- and monoglycerides; phospholipids such as phosphatidyl-choline or phosphatidyl-ethanolamine; or any other pharmaceutically acceptable combined form.

The fatty acids are not toxic and so they may be given in doses of from 1 mg to 100 g per day, preferably 20 mg to 10 g and very preferably 50 mg to 2g/day. They may be administered orally, enterally, parenterally or topically by any appropriate formulation including capsules, pastilles, tablets, powders, emulsions, suspensions, oils, creams, lotions, patches, liposomes, galactosomes or any other form known to those skilled in the art.

The use of DHA as such is preferred. The ratio of DHA to n-6 acids, when present, in the formulations may range from 1:100 to 100:1, preferably 5:1 to 1:5 and very preferably 3:1 to 1:3.

Examples

The following formulations and their use illustrate the invention, by way of example.

- 1. For use in combating dyslexia, soft gelatin capsules or hard gelatin capsules, or pastilles or tablets or other pharmaceutical or nutritional dosage forms containing 100 mg DHA and optionally 100 mg GLA, 100 mg AA and/or 100 mg EPA, for pre-emptive consumption by women in pregnancy or by infants, or for consumption by diagnosed dyslexic children or adults, to give a daily dose of DHA of 20 mg to 10 g.
- Granules or powder for use as above, made with gum acacia, gelatin, starch or other appropriate material containing by weight in each gram, 50 mg DHA, optionally with 50 mg of DGLA, 50 mg AA and/or 50 mg SA.
- 3. Oils for use as above for use as salad oils or for incorporation into any appropriate food material containing 5% by weight DHA, optionally with 10% by weight GLA, 5% by weight AA and 5% by weight EPA.

WO 96/37200 PCT/GB96/01265 5


- 4. Whips, foams, creams, mousses or other liquid or semi-liquid formulations for use as above as foods and containing 2% DHA and optionally 2% by weight GLA, 2% AA, 2% DGLA, 2% SA and 2% EPA.
- 5. Creams, ointments, lotions, shampoos, patches, sticks, pessaries, suppositories or any other dosage form for use as above for topical application in which the active material is an oil containing 3% DHA, optionally with 5% by weight DGLA, 2% AA and 3% EPA.
- 6-10. Formulations as in 1-5 but in which the active ingredients are DHA with any one (or more than one) of the n-6 EFAs selected from GLA, DGLA and AA, and/or any one (or more than one) of the n-3 EFAs selected from SA, 20:4n-3, DPA or EPA.
- 11-15. Formulations as in 1-5 but in which the active ingredients are DHA and either GLA or DGLA of the n-6 series and/or EPA of the n-3 series.
- 16-20. Formulations as in 1-5 but for use in improvement of dark adaptation and/or nocturnal vision generally in dyslexic or non-dyslexic individuals.

CLAIMS

- A method of combating dyslexia, or inadequate night vision or dark adaptation in dyslexics or normal persons, and a method of manufacture of a medicament for such purposes, characterised by administering/incorporating DHA as an active in effective amount.
- A method according to Claim 1, administration being to children or adults showing one or more of the conditions.
- 3. A preventative method according to Claim 1, administration being to women during pregnancy or to infants in the year after birth.
- 4. A method according to any preceding claim wherein one or more n-6 EFAs drawn from the group consisting of linoleic, gamma-linolenic, dihomogamma-linolenic and arachidonic acids and/or one or more n-3 EFAs drawn from the group consisting of alpha-linolenic, stearidonic and eicosapentaenoic acids is used in addition to the DHA.
- 5. A method according to Claim 1, 2 or 3 wherein a precursor n-3 essential fatty acid is substituted for all or part of the DHA.
- 6. A method according to Claim 1, 2 or 3 wherein administration is of, or the medicament is suited to administration of, 1 mg to 100 g, preferably 20 mg to 10 g and very preferably 50 mg to 2 g per day of the DHA.
- 7. A method according to Claim 4 or 5 wherein administration is of, or the medicament is suited to administration of, amounts of fatty acids that in total are within the ranges in Claim 6.
- 8. A method according to Claim 4 wherein administration is of, or the medicament is suited to administration of, DHA and n-6 essential fatty acids in a weight ratio of 1:100 to 100:1, preferably 5:1 to 1:5 and very preferably 3:1 to 1:3.
- 9. A method according to any preceding claim wherein the DHA and/or other essential fatty acids are present as one or more of:- salts including lithium salts; esters; diesters of 1,3-propane diol and other diols; amides; fatty acid alcohols; mono-, di- or

SUBSTITUTE SHEET (RULE 26)


triglycerides; phosphatidyl-choline, phosphatidyl-ethanolamine or other phospholipid; or other pharmaceutically acceptable combined form.



SUBSTITUTE SHEET (RULE 26)

Fig. 2


Figure 2 Dark adaptation in dyslexics compared with non-dyslexic controls



SUBSTITUTE SHEET (RULE 26)

Fig. 3

Figure 3 Effect of Fish Oil Supplementation on Dark Adaptation in Dyslexics



INTERNATIONAL SEARCH REPORT

Inter onal Application No PCT/GB 96/01265

			
A. CLASSI IPC 6	A61K31/20 //(A61K31/20,A61K31:2	9)	
	•		
According to	o International Patent Classification (IPC) or to both national classifi	cation and IPC	
	SEARCHED		
IPC 6	ocumentation searched (classification system followed by classification A61K	on symbols)	
Documentat	tion searched other than minimum documentation to the extent that st	uch documents are included in the fields s	earched
Electronic d	ata base consulted during the international search (name of data base	and, where practical, search terms used)	
	AND CONTROL OF THE PARTY AND		
	TENTS CONSIDERED TO BE RELEVANT	levent nerverer	Relevant to claim No.
Category *	Citation of document, with indication, where appropriate, of the rel	levant passages	Nete value to classify the
X.	LIPIDS,		1-9
	vol. 29, no. 1, - 1994 USA, pages 61-65, XP000600054		
	HOLMAN ET AL: "ABNORMAL PLASMA L	IPIDS OF	
·	PATIENTS WITH RETINITS PIGMENTOSA		
	see abstract		
P.Y	MEDICAL HYPOTHESIS.		1-9
'''	vol. 45, no. 6, - 1995 USA,		
	pages 605-613, XP000600048		
ł	HORROBIN ET AL : "POSSIBLE RELEV PHOSPHOLIPID ABNORMALITIES AND GE		
	INTERACTIONS IN PSYCHIATRIC DISC		
·	THE RELATIONSHIP BETWEEN DYSLEXIA		
	SCHIZOPHRENIA"	•	•
ĺ	see the whole document		
!	-	/	
		•	
X Fun	ther documents are listed in the continuation of hox C.	X Patent family members are listed	in annex.
* Special ca	ategories of cited documents:	T" later document published after the int	ernational filing date
'A' docum	nent defining the general state of the art which is not	or priority date and not in conflict w cited to understand the principle or t	heory underlying the
considered to be of particular relevance invention E earlier document but published on or after the international 'X' document of particular relevance; the claimed invention			claimed invention
filing date cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone.			ocument is taken alone
which is cited to establish the publication date of another 'Y' document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the			eventive step when the
O' document referring to an oral disclosure, use, exhibition or document is combined with one or more other such documents, such combination being obvious to a person skilled			
"P" docum	'P' document published prior to the international filing date but in the art. later than the priority date claimed '&' document member of the same patent family		
Date of the	e actual completion of the international search	Date of mailing of the international s	earch report
g	9 September 1996	24.09.96	
Name and	mailing address of the ISA European Patent Office, P.B. 5818 Patentlaan 2	Authorized officer	
	NL - 2280 HV Rijswijk Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,	Herrera. S	
1	21 701 740 7014	i nerrera. 3	•

Form PCT/ISA/210 (second sheet) (July 1992

INTERNATIONAL SEARCH REPORT

Intes Jonal Application No PCT/GB 96/01265

C.(Continuati	on) DOCUMENTS CONSIDERED TO BE RELEVANT	
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,Y	LANCET, vol. 345, no. 8963, - 10 June 1995 pages 1463-1468, XP000579670 M.MAKRIDES ET AL: "ARE LONG-CHIN POLYUNSATURATED FATTY ACIDS ESSENTIAL NUTRIENTS IN INFANCY?" see the whole document	1-3
Y	EP,A,O 599 576 (SCOTIA HOLDINGS PLC) 1 June 1994 see the whole document	1-9
P,Y	EP,A,O 711 503 (SCOTIA HOLDINGS PLC) 15 May 1996 see the whole document	1-9
·		

1

INTERNATIONAL SEARCH REPORT Inte. Jonal Application No

Information on patent family members

Inte. Jonal Application No PCT/GB 96/01265

Patent document cited in search report	Publication date	Patent memb		Publication date
EP-A-0599576	01-06-94	AU-B- AU-B- AU-B- CA-A- CN-A- JP-A- NO-A- US-A- ZA-A-	666782 5183093 5232996 2109777 1104494 6199663 934266 5516800 9308835	22-02-96 09-06-94 18-07-96 27-05-94 05-07-95 19-07-94 27-05-94 14-05-96 02-08-94
EP-A-0711503	15-05-96	AU-B- CA-A- JP-A- NO-A-	3782895 2162739 8205769 954573	23-05-96 15-05-96 13-08-96 15-05-96

This Page is inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:
BLACK BORDERS
MAGE CUT OFF AT TOP, BOTTOM OR SIDES
☐ FADED TEXT OR DRAWING
D BLURED OR ILLEGIBLE TEXT OR DRAWING
O SKEWED/SLANTED IMAGES
O COLORED OR BLACK AND WHITE PHOTOGRAPHS
GRAY SCALE DOCUMENTS
LINES OR MARKS ON ORIGINAL DOCUMENT
☐ REPERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY
OTHER:

IMAGES ARE BEST AVAILABLE COPY.
As rescanning documents will not correct images problems checked, please do not report the problems to the IFW Image Problem Mailbox